

Pseudotumor cerebri presenting with 2nd and 7th nerve palsy

Mathisha Ebby Perin*, Suresh Kanna S, Dinesh Babu

To Cite:

Perin ME, Kanna SS, Babu D. Pseudotumor cerebri presenting with 2nd and 7th nerve palsy. Medical Science 2022; 26:ms351e2295.
doi: <https://doi.org/10.54905/disssi/v26i127/ms351e2295>

Authors' Affiliation:

Department of General Medicine, Sree Balaji Medical College & Hospital, India

*Corresponding Author

Department of General Medicine, Sree Balaji Medical College & Hospital, Chennai, India
Email: ebby.perin@gmail.com

Peer-Review History

Received: 18 May 2022
Reviewed & Revised: 20/May/2022 to 29/August/2022
Accepted: 31 August 2022
Published: 05 September 2022

Peer-review Method

External peer-review was done through double-blind method.

URL: <https://www.discoveryjournals.org/medicalscience>



This work is licensed under a Creative Commons Attribution 4.0 International License.

ABSTRACT

Idiopathic intracranial hypertension, alternatively called pseudotumor cerebri is characterized by increased intracranial pressure of unknown pathology with the absence of any structural lesions. This intracranial hypertension most commonly presents with postural headache, nausea, vomiting, transient visual loss with cranial nerve involvement, with most common VI cranial nerve involvement. In the following case report, we present a pseudotumor cerebri presenting with 2nd and 7th nerve palsy.

Keywords: Hypertension, Nerve Palsy, Pseudotumor Cerebri

1. INTRODUCTION

Idiopathic intracranial hypertension, alternatively called pseudotumor cerebri is a disorder of unknown pathology, which proves any absence of intracranial space occupying lesions or CSF obstruction (Biousse et al., 2012; Albalawi et al., 2021). This IIH is most commonly seen among women of childbearing age mainly in overweight and obese women. It is also been associated with drugs like tetracycline and with OCPs and even with high dose vitamin A derivatives. Patients who are affected with IIH 92% present with headache, 72% with visual obsurations, 59% with multiple cranial nerve involvement (Peralta & Cestari, 2018). The most commonly involved cranial nerve among them is 6th CN in 12% adults and 40% in children. Very less frequently other nerves are being involved. Very verylimited number of cases has been reported with 2nd and 7th nerve palsy (Acheson, 2006) and even there are even fewer cases reported with 2nd nerve involvement. In the light of this, we present a very rare case with raised ICT involving 2nd nerve and unilateral 7th nerve involvement. This will alert clinicians to suspect IIH in patients with these cranial nerve involvements.

2. CASE REPORT

A 32 year old female came to the ER with the complaints of complaints of headache and giddiness for past 1 week, complaints of blurring of vision for the past 1 week which was gradual in onset, complaints of inability to close right eyes, complaints of diplopia, complaints of deviation of angle of mouth to the left, patient had no complaints of slurring of speech, no complaints of weakness of both upper and lower limbs, no complaints of seizures, vomiting,

trauma. She did not have any long-standing history of headaches, fever or any history of oral contraceptive pills or other significant drug history.

On examination patient is conscious oriented, afebrile, moderately built and nourished with BMI 24. No pallor, icterus, cyanosis, clubbing, pedal edema, lymphadenopathy. Her Vitals: HR: 110/min, BP: 140/90 RR:18/min, Spo2:99% in RA. Systemic examination: CNS- Higher mental function normal, 2nd cranial nerve shows B/L distant 6/18, B/L color vision lost, 3rd, 4th, 6th cranial nerve extra ocular movement full, pupil size b/l 3mm equal, relative afferent pupil defect (RAPD), 7th cranial nerve showed a deviation of angle of mouth to the right side, inability to close left eye, drooling of saliva present, loss of nasolabial fold in left side present all other cranial nerve examinations were absolutely normal, her motor and sensory system were normal, no cerebellar signs, no signs of nystagmus. Other system examination: RS; BAE+ no added sounds, CVS: S1S2+ P/A; soft, no organomegaly.




Figure 1 B/L fundus showing established papilloedema

Fundus examination showed bilateral disc swelling with circumferential hemorrhages (Figure 1). The above findings were confirmed with OCT (Optical Coherence Tomography) of the optic disc. There was severe diffuse thickening of the nerve fibre layer. The thickness of NFL in right and left eye was 240µm and 220µm. Investigations showed CBC (TC:7560, HB: 12, PLT: 2.6 lakh/cumm), ESR:40, CRP:0.6. Her LFT and RFT were within normal limits. Urine routine was normal. Above all, her CT and MRI did not show any significant abnormality.

At day 1 of her admission guarded lumbar puncture was done which showed a opening CSF pressure was more than 35cmH2O and her cerebral fluid analysis showed (protein: 27mg/dl, glucose: 71mg/dl). Therefore a diagnosis of idiopathic intracranial hypertension was made with involvement of 2nd and 7th nerve palsy.

We started the patient on oral steroids (methylprednisolone 40mg BD), we started her on T. acetazolamide 250mg qid and also furosemide was added to her management. Even after putting her on all these drugs and even with guarded lumbar puncture did not improve her symptoms and even her papilloedema, therefore we planned for a the coperitoneal shunt and not for a ventriculo peritoneal shunt because her ventricles were so much dilated in imaging. After the coperitoneal shunt her symptoms subsided with mild residual papilloedema, her visual field defects were normal and patient was discharged.

3. DISCUSSION

We present a case of moderately built 32 year old female with complaints of headache, giddiness, blurring of vision for past 1 week. Here CNS examination showed involvement of second and seventh nerve palsy and her fundus examination showed established papilloedema (Wakerley et al., 2015). Other conditions like lymes disease, sarcoidosis should be ruled out before making a diagnosis of raised ICT because in these conditions also the most common sole nerve to be affected in raised ICT is 7th nerve. All imaging MRI/CT brain should be taken to come to a clear diagnosis of raised ICT (Wall & George, 1991). IIH affects predominantly moderately built to obese individuals with incidence of 11.9 per 100,000 per year.

The pathophysiology of 2nd and 7th nerve palsy in IIH is not yet clearly known and there is no satisfactory hypotheses exist. The extra pressure is known to act on the extra axial facial nerve (Soroken et al., 2016). This theory is very well accepted in seventh and sixth nerve palsy, due to the known fact that sixth nerve having a long course and the most common nerve affected in ICT. But the cause of second nerve palsy getting affected is still a hypothesis. The only supporting hypothesis is that elevation of the

intracranial pressure causes compression of the optic nerve which impairs its axoplasmic flow and causing 2nd nerve palsy and papilloedema (Rangwala & Liu, 2007). There are only very few cases reported with isolated facial nerve palsy and only one or two cases reported with 2nd nerve palsy.

4. CONCLUSION

Idiopathic intracranial hypertension should to always suspect in young obese women, who comes to our OPD with headache and transient loss of vision. The diagnosis of IIH is always being a diagnosis of conclusion other obstructive or structural lesions should be ruled out. Here by our case we would like to notify that raised ICT can also occur with unilateral 7th nerve palsy and very rare presentation with 2nd nerve palsy. The mechanism of this raised ICH is still not clear with 7th and 2nd nerve palsy. And still prompt investigations and various studies are needed to prove its actual hypothesis.

Informed consent

Consent has been obtained from the patient with regards to the use of the pictures for publication.

Funding

This study has not received any external funding.

Conflicts of interest

The authors declare that there are no conflicts of interests.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

1. Acheson JF. Idiopathic intracranial hypertension and visual function. Br Med Bull 2006; 79-80:233-44 doi: 10.1093/bmb/ldl019. PMID: 17242038.
2. Albalawi A, Alharbi A, Altaieb M, Almatrafi M, Alzahuf S, Alanzi O, Alawad A, Saif AM, Almuwarraee S, Alotaibi B, Almishali F, Kassem T, Alrajab F, Almatrafi O, Alshahrani H. Efficacy of Acetazolamide in management of idiopathic intracranial hypertension: A systematic review and meta-analysis. Medical Science 2021; 25(114):1827-1836
3. Biousse V, Bruce BB, Newman NJ, Update on the Pathophysiology and Management of Idiopathic Intracranial Hypertension. J Neurol Neurosurg Psychiat 2012; 83(5):488-94
4. Madriz Peralta G, Cestari DM. An update of idiopathic intracranial hypertension. Curr Opin Ophthalmol 2018; 29(6):495-502 doi: 10.1097/ICU.0000000000000518. PMID: 30169466.
5. Rangwala LM, Liu GT. Pediatric idiopathic intracranial hypertension. Surv Ophthalmol 2007; 52(6):597-617 doi: 10.1016/j.survophthal.2007.08.018. PMID: 18029269.
6. Soroken C, Lacroix L, Korff CM, Combined VIth and VIIth nerve palsy: Consider idiopathic intracranial hypertension. Eur J Paediatr Neurol 2016; 20(2):336-8
7. Wakerley BR, Tan MH, Ting EY. Idiopathic intracranial hypertension. Cephalalgia 2015; 35(3):248-61 doi: 10.1177/0333102414534329. PMID: 24847166.
8. Wall M, George D. Idiopathic intracranial hypertension. A prospective study of 50 patients. Brain 1991; 114 (Pt 1A):155-80 PMID: 1998880.